This product is designed specifically as a general non-acid cleaner and disinfectant for use in hospitals, schools, office buildings and veterinary clinics where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces; floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic file, plastic surfaces, bathrooms, shower stalls, bathrubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical or semi-critical medical device surfaces. Fill or refill the 32 ozs. this product spray bottle with this product refill. Pour product into spray

bottle over sink or tub basis product spray dode with this product refin. Four product fine spray bottle over sink or tub basis or bright or brigh

and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave ort of soap scum.

Disinfection of animal quarters and kennels: Before using this product, remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or trayersed by animals. Empty all troughs, racks, and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Apply this product for disinfection of pre-cleaned animal quarters and kennels. Saturate the surfaces with this product for a period of 3 minutes (for Bovine Viral Diarrhea Virus (BVDV) 5 minutes) (for Feline Calicivrirs and Rabies virus 30 seconds) (for Canine Parvovirus 10 minutes). Vehiliate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse. Using approved ADAC test methods, in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-prous inaminate surfaces: RACTERIDIAN ACTIVITY: Pseudomonas acrunionsa. Listeria

contact time, unless otherwise noted, this product kills the following organisms on hard nonporous inanimate surfaces: BACTERICDIA_ACTIVITY: Pseudomonas aeruginosa, Listeria
monocytogenes, Staphylococcus aureus, Yersina enterocolitica, Salmonella (choleraesuis)
enterica, Enterococcus faecium, Escherichia coii, Corynebacterium armoniagenes,
Escherichia coii 0157:H7, Salmonella (typhi) enterica, Streptococcus pyogenes (Necrotizing
rascritis-Group A) (VA. Medical Center Isolate (AdO)). AMTIBIOTIC-TESISTANT: Methicillin
resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis
(VRE), Vancomycin intermediate, resistant Staphylococcus aureus (VISA), Methicillin
resistant Staphylococcus epidermidis (MRSE). Community Associated Methicillin resistant
Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400, Community Associated
Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300.
TUBERCULOCIDAL ACTIVITY - This product extinibits disinfectant efficacy against
Mycobacterium tuberculosis BCG at 20 degrees Centigrade (20°C) when the treated surface
is allowed to remain wet for 5 minutes.

is allowed to remain wet for 5 minutes.

"Allowed to remain wet for 5 minutes."

"INCLORA ACTIVITY - This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a 10 minute contact time against: Canine Parvovirus, Hepatitis A

Virus (HAV) (University of Ottawa), Poliovirus Type 1 This product, when used on environmental, manimate hard surfaces exhibits virucidal activity against: **Hepatitis B Virus (HBV)**, (Hepadnayrus Testing, against negatilist Virus in 1997 (replandavirus testinistinist), repatrilist C Virus (HCV) and Bovine Viral Diarrhea Virus (80DV) (VIVS.) when the freated surface is allowed to remain wet for 5 minutes; human Coronavirus, SARS associated Coronavirus (ZeptoMetrix) Avian Influenza A strain HSN2 and Avian Influenz Manufactured for:

CARROLLCLEAN

2900 W. Kingsley Rd., • Garland, Texas 75041

800 527 5722 • www carrollclean com

2 minutes; Paramyxovirus (Mumps), Rhinovirus type 39 and Rotavirus (University of Ottawa)

CARROLLCLEAN

OMNI-Q **DETERGENT DISINFECTANT**

EPA Reg. No. 1839-83-4313 EPA Est. No. 4313-TX-1

U.S. PATENT NO. 5.444.094

CLEANER, DEODORIZER, MILDEWSTAT (ON HARD INANIMATE SURFACES), DISINFECTANT, TUBERCULOCIDAL, *VIRUCIDE FUNGICIDE (AGAINST PATHOGENIC FUNGI) KILLS PANDEMIC 2009 H1N1 INFLUENZA A VIRUS (FORMERLY CALLED SWINE FLU)

ACTIVE INGREDIENTS

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈) dimethyl benzyl ammonium chlorides...... 0.105% n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chlorides 0.105%

INERT INGREDIENTS.......99.790% Total 100.000%

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment, if in eyes: Ald eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue insing eye. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting projects and to do so have been control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting projects and to do so have the objects of the control center of doctor. Do unless fold to do so by the poison control center or doctor. Do not you want to the poison control center or doctor. Do not you earlything by mouth to an unconscious person. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible, call a poison control center or doctor for further tréatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may

contraindicate the use of gastric lavage.

KEEP OUT OF REACH OF CHILDREN

See back panels for additional precautionary statements.

> ITEM# 731

for 3 minutes; HIV-1 (associated with AIDS) (Advanced Biotechnologies, Inc.) for 1 minute; and Norovirus (Norwalk virus), Feline Calicivirus (University of Ottawa) and Rabies Virus

(CDC) for 30 seconds. KILLS HIV-1, HBY AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals,

PREVIOUSLY SOILED WITH BLOOD BODY FLUIDS in health care settings (hospitals Nusing Homes) or other settings in which there is an expected likelihood of solid inanimate surfaces objects with blood or body fluids, and in which the surfaces objects likely and the surfaces objects with blood or solid him to potential for transmission of Human Immunodeticlency Virus Type 1 (HIV-1) (associated with AIDS), hepatitis B Virus (HZV), and Hepatitis C Virus HZV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV ON SUFFACES/OBJECTS SOILED WITH BLOOD BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eye coverings. CLEANING PROCEDURES, Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. CONTACT TIME: Allow surface to remain wet for HBV and HCV 5 minutes, for HIV-1 i minute. These contact times will not control other common types of viruses and bacteria. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. FUNGICIOAL ACTIVITY — This product is fungicidal against the pathogenic fungus, Tinchophyton mentagrophytes (Attliete's Foot Fungus) (a cause of ringworm of the foot), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. MILDEWSTRAT — To control mold and mildew on pre-cleaned, fard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew dispressions.

spray sanate to be dealed internal sufe to week online test. Let all diff. He peat application at weekly intervals or when mildew growth appears. Efficacy Tests have demonstrated that this product is an effective Bactericide, fundicide, and virucide in the presence of organic soil (5% BLOOD SERUM). PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

chewing gum or using tobacco.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL

PESTICIDE STORAGE Store in a dry place no lower in temperature than 50°F or higher than 120°F. PESTICIDE DISPOSAL. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or insale is a violation of Federal Law. It these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL — Nonrefillable container, Do not reuse or refill container. Clean container promptly after emptying, Triple rinse as follows: Fill container ½, full with water and recap, Aditate vigorously. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available puncture and dispose in a sanitary landfill. CONTAINER DISPOSAL FOR REFILL SPRAY BOTTLE: Do not reuse empty spray bottle except with empty refill bottle. Wrao

refill bottle, otherwise discard in trash. Do not reuse refill bottle and discard in trash. Offer for recycling or reconditioning, if available.

empty refill bottle. Wrap

731-20130410